


FBXO43 (Human) Recombinant Protein (Q01)

Catalog # H00286151-Q01

Size 10 ug, 25 ug

Applications



Specification

Product Description	Human FBXO43 partial ORF (XP_209918, 1 a.a. - 110 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MSQRHSGQAGTEAGNGADSPPIVNSKYSTRDFCSTSSFQDSGYNELKSCSFDNIDKEYLGKKE KGPTLLYEHPEPESGLGLTHPLESPTQKKKCILPRKEKDKTPELCET
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	37.84
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — FBXO43

Entrez GeneID [286151](#)

GeneBank Accession# [XM_209918](#)

Protein Accession# [XP_209918](#)

Gene Name FBXO43

Gene Alias EMI2, ERP1, Fbx43

Gene Description F-box protein 43

Omim ID [609110](#)

Gene Ontology [Hyperlink](#)

Gene Summary Members of the F-box protein family, such as FBXO43, are characterized by an approximately 40 -amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM]

Other Designations early mitotic inhibitor 2