

Histone H3.1 (Human) Cell-Based ELISA Kit

Catalog # KA2761

Size 1 Kit

Specification

Product Description	Histone H3.1 (Human) Cell-Based ELISA Kit is an indirect enzyme-linked immunoassay for qualitative determination of Histone H3 expression in cultured cells.
Suitable Sample	Attached Cell, Loosely Attached Cell, Suspension Cell
Label	HRP-conjugated
Detection Method	Colorimetric
Assay Type	Qualitative
Reactivity	Human, Mouse, Rat
Regulation Status	For research use only (RUO)
Storage Instruction	Store the kit at 4°C.

Applications

- Qualitative

Gene Info — HIST1H3A

Entrez GeneID	8350
Protein Accession#	P68431
Gene Name	HIST1H3A
Gene Alias	H3/A, H3FA
Gene Description	histone cluster 1, H3a

Omim ID	602810
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]
Other Designations	H3 histone family, member A histone 1, H3a

Gene Info — HIST1H3D

Entrez GeneID	8351
Protein Accession#	P68431
Gene Name	HIST1H3D
Gene Alias	H3/b, H3FB
Gene Description	histone cluster 1, H3d
Omim ID	602811
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]
Other Designations	H3 histone family, member B OTTHUMP00000016149 histone 1, H3d

Gene Info — HIST1H3C

Entrez GeneID	8352
Protein Accession#	P68431

Gene Name	HIST1H3C
Gene Alias	H3.1, H3/c, H3FC
Gene Description	histone cluster 1, H3c
Omim ID	602812
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]
Other Designations	H3 histone family, member C histone 1, H3c

Gene Info — HIST1H3E

Entrez GeneID	8353
Protein Accession#	P68431
Gene Name	HIST1H3E
Gene Alias	H3.1, H3/d, H3FD
Gene Description	histone cluster 1, H3e
Omim ID	602813
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]
Other Designations	H3 histone family, member D histone 1, H3e

Gene Info — HIST1H3I

Entrez GeneID [8354](#)**Protein Accession#** [P68431](#)**Gene Name** HIST1H3I**Gene Alias** H3.f, H3/f, H3FF**Gene Description** histone cluster 1, H3i**Omim ID** [602814](#)**Gene Ontology** [Hyperlink](#)

Gene Summary

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]

Other Designations H3 histone family, member F|OTTHUMP00000017803|histone 1, H3i

Gene Info — HIST1H3G

Entrez GeneID [8355](#)**Protein Accession#** [P68431](#)**Gene Name** HIST1H3G**Gene Alias** H3/h, H3FH**Gene Description** histone cluster 1, H3g**Omim ID** [602815](#)**Gene Ontology** [Hyperlink](#)

Gene Summary

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]

Other Designations

H3 histone family, member H|OTTHUMP00000016152|histone 1, H3g

Gene Info — HIST1H3J**Entrez GeneID**

[8356](#)

Protein Accession#

[P68431](#)

Gene Name

HIST1H3J

Gene Alias

H3/j, H3FJ

Gene Description

histone cluster 1, H3j

Omim ID

[602817](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]

Other Designations

H3 histone family, member J|OTTHUMP00000017804|histone 1, H3j

Gene Info — HIST1H3H**Entrez GeneID**

[8357](#)

Protein Accession#

[P68431](#)

Gene Name

HIST1H3H

Gene Alias

FLJ92264, H3/k, H3F1K, H3FK

Gene Description	histone cluster 1, H3h
Omim ID	602818
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]
Other Designations	H3 histone family, member K histone 1, H3h

Gene Info — HIST1H3B

Entrez GeneID	8358
Protein Accession#	P68431
Gene Name	HIST1H3B
Gene Alias	H3/I, H3FL
Gene Description	histone cluster 1, H3b
Omim ID	602819
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]
Other Designations	H3 histone family, member L OTTHUMP00000016132 histone 1, H3b

Gene Info — HIST1H3F

Entrez GeneID	8968
---------------	----------------------

Protein Accession#	P68431
Gene Name	HIST1H3F
Gene Alias	H3/i, H3FI
Gene Description	histone cluster 1, H3f
Omim ID	602816
Gene Ontology	Hyperlink
Gene Summary	<p>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq]</p>
Other Designations	H3 histone family, member OTTHUMP00000016151 histone 1, H3f

Pathway

- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)
- [Systemic lupus erythematosus](#)

Disease

- [Abortion](#)
- [Genetic Predisposition to Disease](#)
- [Tobacco Use Disorder](#)