


# C1QTNF2 HEK293 Cell Transient Overexpression Lysate(Non-Denatured)


Catalog # L030T6      Size 100 ug

## Applications



### SDS-PAGE Gel

C1QTNF2 transfected lysate



### Western Blot

Lane 1: C1QTNF2 transfected lysate ( 30 KDa).

Lane 2: Non-transfected lysate.

## Specification

Transfected Cell Line	HEK293
Plasmid	pCMV-C1QTNF2 full length
Host	Human
Theoretical MW (kDa)	30
Lysis Buffer	Modified RIPA Lysis Buffer:50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Concentration	2 mg/ml

**Quality Control Testing**

Transient overexpression cell lysate was tested with Anti-C1QTNF2 antibody ([H00114898-M01](#)) by Western Blots.  
SDS-PAGE Gel  
C1QTNF2 transfected lysate  
Western Blot  
Lane 1: C1QTNF2 transfected lysate ( 30 KDa).  
Lane 2: Non-transfected lysate.

**Recommend Usage**

Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.

**Storage Buffer**

In modified RIPA Lysis Buffer.

**Storage Instruction**

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot
- Immunoprecipitation

[Protocol Download](#)

## Gene Info — C1QTNF2

**Entrez GeneID**

[114898](#)

**GeneBank Accession#**

[BC011699](#)

**Protein Accession#**

[AAH11699](#)

**Gene Name**

C1QTNF2

**Gene Alias**

CTRP2, zacrp2

**Gene Description**

C1q and tumor necrosis factor related protein 2

**Gene Ontology**

[Hyperlink](#)

**Other Designations**

complement-c1q tumor necrosis factor-related protein 2