


C20orf102 HEK293 Cell Transient Overexpression Lysate(Non-Denatured)


Catalog # L040T6 Size 100 ug

Applications



SDS-PAGE Gel

C20orf102 transfected lysate



Western Blot

Lane 1: C20orf102 transfected lysate (22 KDa).

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	HEK293
Plasmid	pCMV-C20orf102 full length
Host	Human
Theoretical MW (kDa)	22
Lysis Buffer	Modified RIPA Lysis Buffer:50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Concentration	2 mg/ml

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-C20orf102 antibody ([H00128434-M01](#)) by Western Blots.
SDS-PAGE Gel
C20orf102 transfected lysate
Western Blot
Lane 1: C20orf102 transfected lysate (22 KDa).
Lane 2: Non-transfected lysate.

Recommend Usage

Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Storage Buffer

In modified RIPA Lysis Buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunoprecipitation

[Protocol Download](#)

Gene Info — VSTM2L

Entrez GeneID [128434](#)

GeneBank Accession# [BC033818](#)

Protein Accession# [AAH33818](#)

Gene Name VSTM2L

Gene Alias C20orf102, dJ1118M15.2

Gene Description V-set and transmembrane domain containing 2 like

Gene Ontology [Hyperlink](#)

Other Designations OTTHUMP00000030945