


G22P1 HEK293 Cell Transient Overexpression Lysate(Non-Denatured)

Catalog # L105T6


Size 100 ug

Applications



SDS-PAGE Gel

XRCC6 transfected lysate



Western Blot

Lane 1: XRCC6 transfected lysate (70 KDa).

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line HEK293

Plasmid pCMV-G22P1 full length

Host Human

Theoretical MW (kDa) 70

Lysis Buffer Modified RIPA Lysis Buffer:50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.

Concentration 2 mg/ml

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-G22P1 antibody ([H00002547-M01](#)) by Western Blots.
SDS-PAGE Gel
XRCC6 transfected lysate
Western Blot
Lane 1: XRCC6 transfected lysate (70 KDa).
Lane 2: Non-transfected lysate.

Recommend Usage

Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Storage Buffer

In modified RIPA Lysis Buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunoprecipitation

[Protocol Download](#)

Gene Info — XRCC6

Entrez GeneID [2547](#)

GeneBank Accession# [BC008343](#)

Protein Accession# [AAH08343](#)

Gene Name XRCC6

Gene Alias CTC75, CTCBF, G22P1, KU70, ML8, TLAA

Gene Description X-ray repair complementing defective repair in Chinese hamster cells 6

Omim ID [152690](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq]

Other Designations

ATP-dependent DNA helicase II, 70 kDa subunit|CTC box binding factor 75 kDa subunit|Ku autoantigen p70 subunit|Ku autoantigen, 70kDa|OTTHUMP00000028581|thyroid autoantigen 70kD (Ku antigen)|thyroid autoantigen 70kDa (Ku antigen)|thyroid-lupus autoantigen

Pathway

- [Non-homologous end-joining](#)

Disease

- [Adenocarcinoma](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Central Nervous System Neoplasms](#)
- [Diabetes Mellitus](#)
- [DNA Damage](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)

- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Male Urogenital Diseases](#)
- [Micronuclei](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Pterygium](#)
- [Radiodermatitis](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)