


MAP3K5 HEK293 Cell Transient Overexpression Lysate(Non-Denatured)

Catalog # L141T6


Size 100 ug

Applications



SDS-PAGE Gel

MAP3K5 transfected lysate



Western Blot

Lane 1: MAP3K5 transfected lysate (155 KDa).

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line	HEK293
Plasmid	pCMV-MAP3K5 full length
Host	Human
Theoretical MW (kDa)	155
Lysis Buffer	Modified RIPA Lysis Buffer:50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Concentration	2 mg/ml

Quality Control Testing

Transient overexpression cell lysate was tested with Anti-MAP3K5 antibody ([H00004217-M03](#)) by Western Blots.
SDS-PAGE Gel
MAP3K5 transfected lysate
Western Blot
Lane 1: MAP3K5 transfected lysate (155 KDa).
Lane 2: Non-transfected lysate.

Recommend Usage

Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Storage Buffer

In modified RIPA Lysis Buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot
- Immunoprecipitation

[Protocol Download](#)

Gene Info — MAP3K5

Entrez GeneID [4217](#)

GeneBank Accession# [BC054503](#)

Protein Accession# [AAH54503](#)

Gene Name MAP3K5

Gene Alias ASK1, MAPKKK5, MEKK5

Gene Description mitogen-activated protein kinase kinase kinase 5

Omim ID [602448](#)

Gene Ontology [Hyperlink](#)

Gene Summary

Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK. [provided by RefSeq]

Other Designations

MAP/ERK kinase kinase 5|MAPK/ERK kinase kinase 5|OTTHUMP00000017275|apoptosis signal regulating kinase

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)

Disease

- [Asthma](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Insulin Resistance](#)
- [Lymphoma](#)
- [Tobacco Use Disorder](#)