


FGF2 monoclonal antibody, clone 2H5G2C11


Catalog # MAB10647 Size 100 uL

Applications



Western Blot (Recombinant protein)

Western blot analysis using FGF2 monoclonal antibody, clone 2H5G2C11 (Cat # MAB10647) against truncated FGF2 recombinant protein.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human rectum adenocarcinoma tissue showing cytoplasmic localization using FGF2 monoclonal antibody, clone 2H5G2C11 (Cat # MAB10647) with DAB staining.

Specification

Product Description	Mouse monoclonal antibody raised against partial recombinant FGF2.
Immunogen	Recombinant protein corresponding to human FGF2.
Host	Mouse
Reactivity	Human
Form	Liquid

Recommend Usage	ELISA (1:10000) Western Blot (1:500-1:2000) Immunohistochemistry (1:200-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In ascites (0.03% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Recombinant protein)

Western blot analysis using FGF2 monoclonal antibody, clone 2H5G2C11 (Cat # MAB10647) against truncated FGF2 recombinant protein.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human rectum adenocarcinoma tissue showing cytoplasmic localization using FGF2 monoclonal antibody, clone 2H5G2C11 (Cat # MAB10647) with DAB staining.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — FGF2

Entrez GeneID	2247
Gene Name	FGF2
Gene Alias	BFGF, FGFB, HBGF-2
Gene Description	fibroblast growth factor 2 (basic)
Omim ID	134920
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. [provided by RefSeq]

Other Designations

basic fibroblast growth factor bFGF|fibroblast growth factor 2|heparin-binding growth factor 2|prostatropin

Pathway

- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Alzheimer disease](#)
- [Arthritis](#)
- [Birth Weight](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Depressive Disorder](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetic Retinopathy](#)
- [Edema](#)

- [Fetal Membranes](#)
- [Gastrointestinal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Metabolic Syndrome X](#)
- [Myocardial Infarction](#)
- [Myopia](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)