


BAX monoclonal antibody, clone SPM336

Catalog # MAB12019 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma with BAX monoclonal antibody, clone SPM336 (Cat # MAB12019).

Specification

Product Description	Mouse monoclonal antibody raised against synthetic peptide of BAX.
Immunogen	A synthetic peptide corresponding to amino acids 3-16 of human BAX.
Sequence	GSGEQPRGGGPTSS
Host	Mouse
Theoretical MW (kDa)	21
Reactivity	Human
Specificity	Does not react with mouse and rat.
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1
Recommend Usage	Immunohistochemistry (0.5-1 ug/mL) Western Blot (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.

Storage Buffer	In 10mM PBS (0.05% BSA, 0.05% sodium azide).
Storage Instruction	Store at 4°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma with BAX monoclonal antibody, clone SPM336 (Cat # MAB12019).

Gene Info — BAX

Entrez GeneID	581
Gene Name	BAX
Gene Alias	BCL2L4
Gene Description	BCL2-associated X protein
Omim ID	600040
Gene Ontology	Hyperlink

Gene Summary	The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq]
Other Designations	apoptosis regulator BAX

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Neurotrophin signaling pathway](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)
- [Prion diseases](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia Telangiectasia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Colon cancer](#)
- [Colonic Neoplasms](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Head and Neck Neoplasms](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)

- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Occupational Diseases](#)
- [Osteomyelitis](#)
- [Parkinson disease](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)