


BCL2L1 monoclonal antibody, clone SPM165


Catalog # MAB12023 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of BCL2L1 in Jurkat cell lysate using BCL2L1 monoclonal antibody, clone SPM165 (Cat # MAB12023).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma with BCL2L1 monoclonal antibody, clone SPM165 (Cat # MAB12023).

Specification

Product Description	Mouse monoclonal antibody raised against synthetic peptide of BCL2L1.
Immunogen	A synthetic peptide corresponding to amino acids 33-50 of mouse BCL2L1.
Sequence	NRTEAPEGTESEMETPSA
Host	Mouse
Theoretical MW (kDa)	27
Reactivity	Human
Form	Liquid

Purification	Protein A/G purification
Isotype	IgG2a
Recommend Usage	Immunohistochemistry (0.5-1 ug/mL) Western Blot (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10mM PBS (0.05% BSA, 0.05% sodium azide).
Storage Instruction	Store at 4°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of BCL2L1 in Jurkat cell lysate using BCL2L1 monoclonal antibody, clone SPM165 (Cat # MAB12023).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma with BCL2L1 monoclonal antibody, clone SPM165 (Cat # MAB12023).

Gene Info — BCL2L1

Entrez GeneID	598
Gene Name	BCL2L1
Gene Alias	BCL-XL/S, BCL2L, BCLX, Bcl-X, DKFZp781P2092, bcl-xL, bcl-xS
Gene Description	BCL2-like 1
Omim ID	600039
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Two alternatively spliced transcript variants, which encode distinct isoforms, have been reported. The longer isoform acts as an apoptotic inhibitor and the shorter form acts as an apoptotic activator. [provided by RefSeq]

Other Designations

OTTHUMP00000030550|OTTHUMP00000030551|OTTHUMP00000030553

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Chronic myeloid leukemia](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer Disease](#)
- [Amnesia](#)
- [Cognition Disorders](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)

- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [Neoplasm Metastasis](#)
- [Neuropsychological Tests](#)
- [Occupational Diseases](#)
- [Ovarian Neoplasms](#)
- [Prostatic Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)