# EPHA5 monoclonal antibody, clone 46CT61.6.4

Catalog # MAB12324 Size 400 uL

# Applications



## Western Blot (Cell lysate)

Western blot analysis of U-251 MG cell line lysates reacted with EPHA5 monoclonal antibody (Cat # MAB12324) at 1:1000 dilution.



### Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining of formalin-fixed and paraffin-embedded human brain tissue reacted with EPHA5 monoclonal antibody (Cat # MAB12324) at 1:10-1:50 dilution.

Specification	
Product Description	Mouse monoclonal antibody raised against partial recombinant human EPHA5.
Immunogen	Recombinant His fusion protein corresponding to human EPHA5.
Host	Mouse
Reactivity	Human
Form	Liquid
Purification	Protein G purification



## **Product Information**

lsotype	lgG1, kappa
Recommend Usage	Immunohistochemistry (1:10-1:50) Western Blot (1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## Applications

• Western Blot (Cell lysate)

Western blot analysis of U-251 MG cell line lysates reacted with EPHA5 monoclonal antibody (Cat # MAB12324) at 1:1000 dilution.

• Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of formalin-fixed and paraffin-embedded human brain tissue reacted with EPHA5 monoclonal antibody (Cat # MAB12324) at 1:10-1:50 dilution.

# Gene Info — EPHA5

Entrez GenelD	<u>2044</u>
Gene Name	EPHA5
Gene Alias	CEK7, EHK1, HEK7, TYRO4
Gene Description	EPH receptor A5
Omim ID	<u>600004</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in th e nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an e xtracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin rece ptors are divided into 2 groups based on the similarity of their extracellular domain sequences an d their affinities for binding ephrin-A and ephrin-B ligands. Two transcript variants encoding differ ent isoforms have been found for this gene.



## **Product Information**

#### **Other Designations**

Eph homology kinase-1|ephrin receptor EphA5|ephrin type-A receptor 5|receptor protein-tyrosine kinase HEK7|tyrosine-protein kinase receptor EHK-1

# Pathway

• Axon guidance

#### Disease

• Tobacco Use Disorder