# CD8 monoclonal antibody, clone CA-8

Catalog # MAB12451 Size 100 ug

## Specification

Product Description	Mouse monoclonal antibody raised against full length native of human CD8.
Immunogen	Native purified CD8 from human thymocytes.
Host	Mouse
Reactivity	Human
Form	Lyophilized
lsotype	lgG2a
Recommend Usage	Immunohistochemistry (1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from ascites fluid (1.2 % sodium acetate, 2 mg BSA, 0.01 mg sodium azide)
Storage Instruction	Store at -20°C. After reconstitution with 1 mL of 1.2% sodium acetate or neutral PBS and concentration will be 100 u g/mL, store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

# Applications

Immunohistochemistry (Frozen sections)

Gene Info — CD8A	
Entrez GenelD	<u>925</u>

😵 Abnova	Product Information
Gene Name	CD8A
Gene Alias	CD8, Leu2, MAL, p32
Gene Description	CD8a molecule
Omim ID	<u>186910 608957</u>
Gene Ontology	Hyperlink
Gene Summary	The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediat es efficient cell-cell interactions within the immune system. The CD8 antigen acts as a corepresso r with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen prese nting cell (APC) in the context of class I MHC molecules. The coreceptor functions as either a hom odimer composed of two alpha chains, or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light cha ins. This gene encodes the CD8 alpha chain isoforms. Multiple transcript variants encoding differ ent isoforms have been found for this gene. [provided by RefSeq
Other Designations	CD8 antigen alpha polypeptide CD8 antigen, alpha polypeptide (p32) Leu2 T-lymphocyte antigen  OKT8 T-cell antigen T cell co-receptor T-cell antigen Leu2 T-cell surface glycoprotein CD8 alpha c hain T-lymphocyte differentiation antigen T8/Leu-2 T8 T-cell ant

Gene Info — CD8B		
<u>926</u>		
CD8B		
CD8B1, LYT3, Leu2, Ly3, MGC119115		
CD8b molecule		
<u>186730</u>		
Hyperlink		
The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediat es efficient cell-cell interactions within the immune system. The CD8 antigen, acting as a corecept or, and the T-cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homod imer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 beta chain isoforms. Multiple alternatively spliced transcript variants encoding distinct membrane associated or secreted isoforms have been described. A pseudoge ne, also located on chromosome 2, has been identified. [provided by RefSeq		
CD8 antigen, beta polypeptide (p37) CD8 antigen, beta polypeptide 1 (p37) CD8b antigen OTTH UMP00000160761 T lymphocyte surface glycoprotein beta chain T-cell surface glycoprotein CD8 beta chain		



## Pathway

- Antigen processing and presentation
- Antigen processing and presentation
- Cell adhesion molecules (CAMs)
- Cell adhesion molecules (CAMs)
- Hematopoietic cell lineage
- Hematopoietic cell lineage
- Primary immunodeficiency
- Primary immunodeficiency
- <u>T cell receptor signaling pathway</u>
- <u>T cell receptor signaling pathway</u>

#### Disease

• Tobacco Use Disorder