


RecomAb™

# Histone H4 (acetyl K8) monoclonal antibody, clone RM201

Catalog # MAB12786


Size 100 ug

## Applications



### Western Blot

Western blot analysis of Lane 1: acid extracts of HeLa cell treated with sodium butyrate, Lane 2: acid extracts of HeLa cell untreated and Lane 3: recombinant Histone H4 with Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat # MAB12786) at 0.5 ug/mL working concentration, showed a band of Histone H4 acetylated at Lysine 8.



### Immunocytochemistry

Immunocytochemical staining of HeLa cells treated with sodium butyrate, using Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat# MAB12786) (red). Actin filaments have been labeled with fluorescein phalloidin (green).

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat# MAB12786) specifically reacts to Histone H4 acetylated at Lysine 8 (K8ac). No cross reactivity with unmodified Lysine 8 (K8 ctrl), acetylated Lysine 5 (K5ac), Lysine 12 (K12ac), Lysine 16 (K16ac), Lysine 20 (K20ac), Lysine 31 (K31ac), or Lysine 91 (K91) in Histone H4.

## Specification

Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H4 (acetyl K8).
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic acetyl peptide corresponding to residues surrounding K8 of human Histone H4.
Sequence	N/A
Specificity	This antibody reacts to Histone H4 acetylated at Lysine 8. No cross reactivity with other acetylated Lysines in Histone H4.
Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	ELISA (0.2 ug/mL-1 ug/mL) Western Blot (0.5 ug/mL-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.

## Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat# MAB12786) specifically reacts to Histone H4 acetylated at Lysine 8 (K8ac). No cross reactivity with unmodified Lysine 8 (K8 ctrl), acetylated Lysine 5 (K5ac), Lysine 12 (K12ac), Lysine 16 (K16ac), Lysine 20 (K20ac), Lysine 31 (K31ac), or Lysine 91 (K91) in Histone H4.

## Applications

### Western Blot

Western blot analysis of Lane 1: acid extracts of HeLa cell treated with sodium butyrate, Lane 2: acid extracts of HeLa cell untreated and Lane 3: recombinant Histone H4 with Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat # MAB12786) at 0.5 ug/mL working concentration, showed a band of Histone H4 acetylated at Lysine 8.

### Immunocytochemistry

Immunocytochemical staining of HeLa cells treated with sodium butyrate, using Histone H4 (acetyl K8) monoclonal antibody, clone RM201 (Cat# MAB12786) (red). Actin filaments have been labeled with fluorescein phalloidin (green).

### Enzyme-linked Immunoabsorbent Assay

## Gene Info — HIST4H4

Entrez GeneID [121504](#)

Protein Accession# [P62805](#)

Gene Name HIST4H4

Gene Alias H4/p, MGC24116

Gene Description histone cluster 4, H4

Gene Ontology [Hyperlink](#)

**Gene Summary**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. [provided by RefSeq]

**Other Designations** histone 4, H4|histone H4

## Pathway

- [Systemic lupus erythematosus](#)