


# CD99 monoclonal antibody, clone MEM-131

Catalog # MAB12936

Size 50 ug

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human thymus with CD99 monoclonal antibody, clone MEM-131 (Cat # MAB12936) at 10 ug/mL working concentration.

## Specification

**Product Description** Mouse monoclonal antibody raised against human CD99.

**Immunogen** Human peripheral blood leukemia T cell line.

**Host** Mouse

**Reactivity** Human

**Form** Liquid

**Purification** Caprylic acid and ammonium sulfate precipitation

**Isotype** IgM

**Recommend Usage**

- Flow Cytometry
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (10 ug/mL)
- Immunoprecipitation
- Western Blot
- The optimal working dilution should be determined by the end user.

**Storage Buffer** In PBS, pH 7.4 (0.09% sodium azide).

**Storage Instruction**

Store at 4°C.  
Aliquot to avoid repeated freezing and thawing.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human thymus with CD99 monoclonal antibody, clone MEM-131 (Cat # MAB12936) at 10 ug/mL working concentration.

- Immunoprecipitation
- Flow Cytometry

## Gene Info — CD99

**Entrez GeneID**[4267](#)**Protein Accession#**[P14209](#)**Gene Name**

CD99

**Gene Alias**

MIC2, MIC2X, MIC2Y

**Gene Description**

CD99 molecule

**Omim ID**[313470 450000](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. Cyclophilin A binds to CD99 and may act as a signaling regulator of CD99. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

**Other Designations**

CD99 antigen|E2 antigen|MIC2 (monoclonal antibody 12E7)|OTTHUMP00000022840|T-cell surface glycoprotein E2|antigen identified by monoclonal 12E7, Y homolog|antigen identified by monoclonal antibodies 12E7, F21 and O13|surface antigen MIC2

## Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [Leukocyte transendothelial migration](#)

## Disease

- [Arthritis](#)
- [Genetic Predisposition to Disease](#)