


B3GAT1 monoclonal antibody, clone NK-1

Catalog # MAB13282 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human spleen with B3GAT1 monoclonal antibody, clone NK-1 (Cat # MAB13282).

Specification	
Product Description	Mouse monoclonal antibody raised against native human B3GAT1.
Immunogen	Human peripheral blood mononuclear cells.
Host	Mouse
Theoretical MW (kDa)	~110
Reactivity	Human
Form	Liquid
Isotype	lgM, kappa
Recommend Usage	Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (2-4 ug/mL for 30 min at R T) (Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA , pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human spleen with B3GAT1 monoclonal antibody, clone NK-1 (Cat # MAB13282).

• Immunofluorescence

Gene Info — B3GAT1	
Entrez GenelD	<u>27087</u>
Protein Accession#	<u>Q9P2W7</u>
Gene Name	B3GAT1
Gene Alias	CD57, GLCATP, GIcAT-P, GIcUAT-P, HNK-1, HNK1, LEU7, NK-1
Gene Description	beta-1,3-glucuronyltransferase 1 (glucuronosyltransferase P)
Omim ID	<u>151290</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the glucuronyltransferase gene family. These en zymes exhibit strict acceptor specificity, recognizing nonreducing terminal sugars and their anom eric linkages. This gene product functions as the key enzyme in a glucuronyl transfer reaction duri ng the biosynthesis of the carbohydrate epitope HNK-1 (human natural killer-1, also known as CD 57 and LEU7). Alternate transcriptional splice variants have been characterized. [provided by Ref Seq
Other Designations	CD57 antigen LEU7 antigen UDP-GlcUA:glycoprotein beta-1,3-glucuronyltransferase beta-1,3-glu curonyltransferase 1 galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1 glucuron osyltransferase P

Pathway

- <u>Chondroitin sulfate biosynthesis</u>
- Heparan sulfate biosynthesis



Product Information

• Metabolic pathways