


ICAM3 monoclonal antibody, clone ICAM3/1019


Catalog # MAB13316 Size 100 ug

Applications



Western Blot (Cell lysate)

Western Blot analysis of HL-60 cell lysate with ICAM3 monoclonal antibody, clone ICAM3/1019 (Cat # MAB13316).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with ICAM3 monoclonal antibody, clone ICAM3/1019 (Cat # MAB13316).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human ICAM3.
Immunogen	Recombinant protein corresponding to full length human ICAM3.
Host	Mouse
Theoretical MW (kDa)	110-160
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification

Isotype	IgG2b, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) Western Blot (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.
Storage Instruction	Store at -20 to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of HL-60 cell lysate with ICAM3 monoclonal antibody, clone ICAM3/1019 (Cat # MAB13316).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with ICAM3 monoclonal antibody, clone ICAM3/1019 (Cat # MAB13316).

- Immunofluorescence

- Flow Cytometry

Gene Info — ICAM3

Entrez GeneID	3385
Protein Accession#	P32942
Gene Name	ICAM3
Gene Alias	CD50, CDW50, ICAM-R
Gene Description	intercellular adhesion molecule 3
Omim ID	146631
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also as a potent signalling molecule. [provided by RefSeq]

Other Designations

intercellular adhesion molecule-3

Pathway

- [Cell adhesion molecules \(CAMs\)](#)

Disease

- [Arthritis](#)
- [Birth Weight](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hepatitis C](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Severe Acute Respiratory Syndrome](#)