


CD99 monoclonal antibody, clone HO36-1.1

Catalog # MAB13350 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Ewing's sarcoma (A) and human testicular carcinoma (B) with CD99 monoclonal antibody, clone HO36-1.1 (Cat # MAB13350).

Specification

Product Description	Mouse monoclonal antibody raised against native human CD99.
Immunogen	Purified E-rosette forming cells from human peripheral blood lymphocytes.
Host	Mouse
Theoretical MW (kDa)	27-32
Reactivity	Human
Specificity	This monoclonal antibody shows a very similar reactivity to other CD99 monoclonal antibodies.
Form	Liquid
Purification	Protein A/G purification

Isotype	IgM, kappa
Recommend Usage	Flow Cytometry (5-10 uL/million cells) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL for 30 min at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA , pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human Ewing's sarcoma (A) and human testicular carcinoma (B) with CD99 monoclonal antibody, clone HO36-1.1 (Cat # MAB13350).

- Immunofluorescence

- Flow Cytometry

Gene Info — CD99

Entrez GeneID	4267
Protein Accession#	P14209
Gene Name	CD99
Gene Alias	MIC2, MIC2X, MIC2Y
Gene Description	CD99 molecule
Omim ID	313470 450000
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. Cyclophilin A binds to CD99 and may act as a signaling regulator of CD99. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

CD99 antigen|E2 antigen|MIC2 (monoclonal antibody 12E7)|OTTHUMP00000022840|T-cell surface glycoprotein E2|antigen identified by monoclonal 12E7, Y homolog|antigen identified by monoclonal antibodies 12E7, F21 and O13|surface antigen MIC2

Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [Leukocyte transendothelial migration](#)

Disease

- [Arthritis](#)
- [Genetic Predisposition to Disease](#)