


BAX monoclonal antibody, clone BAX/962

Catalog # MAB13415 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma (A) and human melanoma (B) with BAX monoclonal antibody, clone BAX/962 (Cat # MAB13415).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human BAX.
Immunogen	Recombinant protein corresponding to full length human BAX.
Host	Mouse
Theoretical MW (kDa)	21
Reactivity	Human
Specificity	This monoclonal antibody is highly specific to Bax and shows no cross-reaction with Bcl-2 or Bcl-X protein.
Form	Liquid
Purification	Protein A/G purification

Isotype	IgG1
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) Western Blot (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% BSA, 0.05% sodium azide).
Storage Instruction	Store at 4°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of Hodgkin's lymphoma (A) and human melanoma (B) with BAX monoclonal antibody, clone BAX/962 (Cat # MAB13415).
- Immunofluorescence
- Flow Cytometry

Gene Info — BAX

Entrez GeneID	581
Protein Accession#	Q07812
Gene Name	BAX
Gene Alias	BCL2L4
Gene Description	BCL2-associated X protein
Omim ID	600040
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq]

Other Designations

apoptosis regulator BAX

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Neurotrophin signaling pathway](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)
- [Prion diseases](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia Telangiectasia](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Colon cancer](#)
- [Colonic Neoplasms](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)

- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genomic Instability](#)
- [Head and Neck Neoplasms](#)
- [Helicobacter Infections](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Occupational Diseases](#)
- [Osteomyelitis](#)
- [Parkinson disease](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)