


CD86 monoclonal antibody, clone C86/1146


Catalog # MAB13462 Size 100 ug

Applications




Western Blot (Cell lysate)

Western Blot analysis of Daudi cell lysate with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human melanoma with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).



Flow Cytometry

Flow cytometric analysis of human PBMCs with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).

Specification

Product Description

Mouse monoclonal antibody raised against full length recombinant human CD86.

Immunogen

Recombinant protein corresponding to full length human CD86.

Host	Mouse
Theoretical MW (kDa)	70
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (2-4 ug/mL) Western Blot (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.
Storage Instruction	Store at -20 to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of Daudi cell lysate with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human melanoma with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).

- Immunofluorescence

- Flow Cytometry

Flow cytometric analysis of human PBMCs with CD86 monoclonal antibody, clone C86/1146 (Cat # MAB13462).

Gene Info — CD86

Entrez GeneID	942
Protein Accession#	P42081
Gene Name	CD86

Gene Alias	B7-2, B70, CD28LG2, LAB72, MGC34413
Gene Description	CD86 molecule
Omim ID	601020
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequences have not been determined. [provided by RefSeq]
Other Designations	B-lymphocyte activation antigen B7-2 B7-2 antigen CD28 antigen ligand 2 CD86 antigen CD86 antigen (CD28 antigen ligand 2, B7-2 antigen) CTLA-4 counter-receptor B7.2

Pathway

- [Allograft rejection](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Acute Disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Bronchiolitis](#)

- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Scleroderma](#)

- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)