


CD44 monoclonal antibody, clone HCAM/1097

Catalog # MAB13474 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with CD44 monoclonal antibody, clone HCAM/1097 (Cat # MAB13474).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human CD44.
Immunogen	Recombinant protein corresponding to full length human CD44.
Host	Mouse
Theoretical MW (kDa)	80-95
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.

Storage Instruction

Store at -20 to -80°C.
Aliquot to avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human tonsil with CD44 monoclonal antibody, clone HCAM/1097 (Cat # MAB13474).

- Immunofluorescence

- Flow Cytometry

Gene Info — CD44

Entrez GeneID [960](#)

Protein Accession# [P16070](#)

Gene Name CD44

Gene Alias CDW44, CSPG8, ECMR-III, HCELL, IN, LHR, MC56, MDU2, MDU3, MGC10468, MIC4, MUTCH-I, Pgp1

Gene Description CD44 molecule (Indian blood group)

Omim ID [107269](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq]

Other Designations CD44 antigen|CD44 antigen (homing function and Indian blood group system)|CDW44 antigen|GP90 lymphocyte homing/adhesion receptor|Hermes antigen|antigen gp90 homing receptor|cell adhesion molecule|cell surface glycoprotein CD44|chondroitin sulfate proteoglycan

Pathway

- [ECM-receptor interaction](#)
- [Hematopoietic cell lineage](#)

Disease

- [Arthritis](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Craniofacial Abnormalities](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Heart Defects](#)
- [Hepatitis B](#)
- [Kidney Failure](#)
- [Mouth Abnormalities](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Ovarian Neoplasms](#)