


CD63 monoclonal antibody, clone LAMP3/968

Catalog # MAB13480 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human prostate carcinoma (A) and human melanoma (B) with CD63 monoclonal antibody, clone LAMP3/968 (Cat # MAB13480).

Specification	
Product Description	Mouse monoclonal antibody raised against full length recombinant human CD63.
Immunogen	Recombinant protein corresponding to full length human CD63.
Host	Mouse
Theoretical MW (kDa)	26-60
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	lgG2a, kappa



Product Information

Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (0.5-1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1-2 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.
Storage Instruction	Store at -20 to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

• Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human prostate carcinoma (A) and human melanoma (B) with CD63 monoclonal antibody, clone LAMP3/968 (Cat # MAB13480).

- Immunofluorescence
- Flow Cytometry

Gene Info — CD63	
Entrez GenelD	<u>967</u>
Protein Accession#	<u>P08962</u>
Gene Name	CD63
Gene Alias	LAMP-3, ME491, MLA1, OMA81H, TSPAN30
Gene Description	CD63 molecule
Omim ID	<u>155740</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known a s the tetraspanin family. Most of these members are cell-surface proteins that are characterized b y the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded prot ein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndro me. Also this gene has been associated with tumor progression. The use of alternate polyadenyla tion sites has been found for this gene. Alternative splicing results in multiple transcript variants en coding different proteins. [provided by RefSeq



Product Information

Other Designations

CD63 antigen|CD63 antigen (melanoma 1 antigen)|granulophysin|lysosome-associated membra ne glycoprotein 3|melanoma 1 antigen|melanoma-associated antigen ME491|melanoma-associated ed antigen MLA1|ocular melanoma-associated antigen|tetraspanin-30

Pathway

• Lysosome