


CHGA monoclonal antibody, clone CHGA/765

Catalog # MAB14231 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human pancreas (A), human parathyroid (B) and human adrenal gland (C) with CHGA monoclonal antibody, clone CHGA/765 (Cat # MAB14231).

Specification

Product Description	Mouse monoclonal antibody raised against full length recombinant human CHGA.
Immunogen	Recombinant protein corresponding to full length human CHGA.
Host	Mouse
Theoretical MW (kDa)	68-75

Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG2a, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 ⁶ cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS (0.05% BSA, 0.05% sodium azide).
Storage Instruction	Store at 4°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human pancreas (A), human parathyroid (B) and human adrenal gland (C) with CHGA monoclonal antibody, clone CHGA/765 (Cat # MAB14231).

- Immunofluorescence

- Flow Cytometry

Gene Info — CHGA

Entrez GeneID	1113
Protein Accession#	P10645
Gene Name	CHGA
Gene Alias	CGA
Gene Description	chromogranin A (parathyroid secretory protein 1)
Omim ID	118910
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. It is found in secretory vesicles of neurons and endocrine cells. This gene product is a precursor to three biologically active peptides; vasostatin, pancreastatin, and parastatin. These peptides act as autocrine or paracrine negative modulators of the neuroendocrine system. Other peptides, including chromostatin, beta-granin, WE-14 and GE-25, are also derived from the full-length protein. However, biological activities for these molecules have not been shown. [provided by RefSeq]

Other Designations

betagranin (N-terminal fragment of chromogranin A)|chromogranin A|parathyroid secretory protein 1

Disease

- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Hypertension](#)
- [Kidney Failure](#)
- [Prostatic Neoplasms](#)
- [Schizophrenia](#)