


# CHGA monoclonal antibody, clone CHGA/765

Catalog # MAB14232      Size 100 ug

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human pancreas (A), human parathyroid (B) and human adrenal gland (C) with CHGA monoclonal antibody, clone CHGA/765 (Cat # MAB14232).

## Specification

<b>Product Description</b>	Mouse monoclonal antibody raised against full length recombinant human CHGA.
<b>Immunogen</b>	Recombinant protein corresponding to full length human CHGA.
<b>Host</b>	Mouse
<b>Theoretical MW (kDa)</b>	68-75

Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG2a, kappa
Recommend Usage	Flow Cytometry (0.5-1 ug/10 <sup>6</sup> cells in 0.1 mL) Immunofluorescence (1-2 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 10 mM PBS.
Storage Instruction	Store at -20 to -80°C. Aliquot to avoid repeated freezing and thawing.

## Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human pancreas (A), human parathyroid (B) and human adrenal gland (C) with CHGA monoclonal antibody, clone CHGA/765 (Cat # MAB14232).

- Immunofluorescence

- Flow Cytometry

## Gene Info — CHGA

Entrez GeneID	<a href="#">1113</a>
Protein Accession#	<a href="#">P10645</a>
Gene Name	CHGA
Gene Alias	CGA
Gene Description	chromogranin A (parathyroid secretory protein 1)
Omim ID	<a href="#">118910</a>
Gene Ontology	<a href="#">Hyperlink</a>

**Gene Summary**

The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. It is found in secretory vesicles of neurons and endocrine cells. This gene product is a precursor to three biologically active peptides; vasostatin, pancreastatin, and parastatin. These peptides act as autocrine or paracrine negative modulators of the neuroendocrine system. Other peptides, including chromostatin, beta-granin, WE-14 and GE-25, are also derived from the full-length protein. However, biological activities for these molecules have not been shown. [provided by RefSeq]

**Other Designations**

betagranin (N-terminal fragment of chromogranin A)|chromogranin A|parathyroid secretory protein 1

**Disease**

- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Hypertension](#)
- [Kidney Failure](#)
- [Prostatic Neoplasms](#)
- [Schizophrenia](#)