


KRT7 monoclonal antibody, clone OV-TL12/30

Catalog # MAB14307 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of ovarian carcinoma (A) and human lung SCC (B) with KRT7 monoclonal antibody, clone OV-TL12/30 (Cat # MAB14307).

Specification

Product Description	Mouse monoclonal antibody raised against native human KRT7.
Immunogen	OTN 11 ovarian carcinoma cell line.
Host	Mouse
Theoretical MW (kDa)	55
Reactivity	Human
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1

Recommend Usage

Flow Cytometry (0.5-1 ug/10⁶ cells in 0.1 mL)
Immunofluorescence (0.5-1 ug/mL)
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer

In 10 mM PBS (0.05% BSA, 0.05% sodium azide).

Storage Instruction

Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of ovarian carcinoma (A) and human lung SCC (B) with KRT7 monoclonal antibody, clone OV-TL12/30 (Cat # MAB14307).

- Immunofluorescence

- Flow Cytometry

Gene Info — KRT7

Entrez GeneID

[3855](#)

Protein Accession#

[P08729](#)

Gene Name

KRT7

Gene Alias

CK7, K2C7, K7, MGC129731, MGC3625, SCL

Gene Description

keratin 7

Omim ID

[148059](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels. The genes encoding the type II cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing may result in several transcript variants; however, not all variants have been fully described. [provided by RefSeq]

Other Designations

cytokeratin 7|keratin, 55K type II cytoskeletal|keratin, simple epithelial type I, K7|keratin, type II cytoskeletal 7|sarcolectin|type II mesothelial keratin K7
