


# LAMC1 monoclonal antibody, clone CL3195


Catalog # MAB15794      Size 100 uL

## Applications



### Western Blot (Recombinant protein)

Western Blot analysis of recombinant protein Lane 1: Laminin-111, Lane 2: Laminin-121, Lane 3: Laminin-211, Lane 4: Laminin-221, Lane 5: Laminin-411, Lane 6: Laminin-421, Lane 7: Laminin-511, Lane 8: Laminin-521 and Lane 9: Laminin-332 with LAMC1 monoclonal antibody, clone CL3195 (Cat # MAB15794).



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human rectum with LAMC1 monoclonal antibody, clone CL3195 (Cat # MAB15794) shows strong immunoreactivity in basement membrane of glandular epithelium.

## Specification

<b>Product Description</b>	Mouse monoclonal antibody raised against partial recombinant human LAMC1.
<b>Immunogen</b>	Recombinant protein corresponding to human LAMC1.
<b>Sequence</b>	ECREGFVGNRCDQCEENFYNRSWPGCQECPCYRLVKDKVADHRVKLQELESLIANLGTGDE MVTDAQAFEDRLKEAEREVMDLLREAQDVKDQNLMDRLQRVNNTLSSQISRLQNIRNTIEETGN LAEQARAHVENTERLIEIASR
<b>Host</b>	Mouse
<b>Reactivity</b>	Human

Form	Liquid
Purification	Protein A purification
Isotype	IgG2b
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:500-1:1000) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot (Recombinant protein)

Western Blot analysis of recombinant protein Lane 1: Laminin-111, Lane 2: Laminin-121, Lane 3: Laminin-211, Lane 4: Laminin-221, Lane 5: Laminin-411, Lane 6: Laminin-421, Lane 7: Laminin-511, Lane 8: Laminin-521 and Lane 9: Laminin-332 with LAMC1 monoclonal antibody, clone CL3195 (Cat # MAB15794).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human rectum with LAMC1 monoclonal antibody, clone CL3195 (Cat # MAB15794) shows strong immunoreactivity in basement membrane of glandular epithelium.

## Gene Info — LAMC1

Entrez GeneID	<a href="#">3915</a>
Protein Accession#	<a href="#">P11047</a>
Gene Name	LAMC1
Gene Alias	LAMB2, MGC87297
Gene Description	laminin, gamma 1 (formerly LAMB2)
Omim ID	<a href="#">150290</a> <a href="#">176780</a>
Gene Ontology	<a href="#">Hyperlink</a>

## Gene Summary

Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins are composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively) and they form a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isomers combine to give rise to different heterotrimeric laminin isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological functions of the different chains and trimer molecules are largely unknown, but some of the chains have been shown to differ with respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the gamma chain isoform laminin, gamma 1. The gamma 1 chain, formerly thought to be a beta chain, contains structural domains similar to beta chains, however, lacks the short alpha region separating domains Ia and II. The structural organization of this gene also suggested that it had diverged considerably from the beta chain genes. Embryos of transgenic mice in which both alleles of the gamma 1 chain gene were inactivated by homologous recombination, lacked basement membranes, indicating that laminin, gamma 1 chain is necessary for laminin heterotrimer assembly. It has been inferred by analogy with the strikingly similar 3' UTR sequence in mouse laminin gamma 1 cDNA, that multiple polyadenylation sites are utilized in human to generate the 2 different sized mRNAs (5.5 and 7.5 kb) seen on Northern analysis. [provided by RefSeq]

## Other Designations

OTTHUMP00000033450|formerly LAMB2|laminin, gamma 1

## Pathway

- [ECM-receptor interaction](#)
- [Focal adhesion](#)
- [Pathways in cancer](#)
- [Prion diseases](#)
- [Small cell lung cancer](#)

## Disease

- [Genetic Predisposition to Disease](#)
- [Macular Degeneration](#)
- [Ovarian Neoplasms](#)