

EP300 monoclonal antibody

Catalog # MAB15840 Size 50 ug

Applications





ChIP

ChIP was performed using HeLa cells. A titration of the antibody consisting of 1, 2, 5 and 10 ug per ChIP experiment was analysed. IgG (2 ug/IP) was used as negative IP control. Quantitative PCR was performed with primers for two genomic regions near the ANKRD32 and IRS2 genes, used as positive controls, and for the coding region of the inactive MYOD1 gene and an intergeic region on chromosome 11, used as negative controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).

ChIP-Seq

ChIP was performed with 5 ug of antibody. The figure shows the peak distribution along the complete sequence and a 3 mb region of chromosome 5 and in two regions surrounding the IRS2 and ANKRD32 positive control genes. The position of the amplicon used for ChIP-qPCR is indicated by an arrow.

Specification	
Product Description	Mouse monoclonal antibody raised against EP300.
Immunogen	Human p300 by DNA immunization in which the C-terminus of the protein was cloned and expressed .
Host	Mouse
Reactivity	Human
Form	Liquid

😵 Abnova

Product Information

Purification	Protein A purification
lsotype	lgG3
Recommend Usage	ChIP (5 ug/CHIP) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide, 0.05% proclin 300).
Storage Instruction	Store at -20°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

ChIP

ChIP was performed using HeLa cells. A titration of the antibody consisting of 1, 2, 5 and 10 ug per ChIP experiment was analysed. IgG (2 ug/IP) was used as negative IP control. Quantitative PCR was performed with primers for two genomic regions near the ANKRD32 and IRS2 genes, used as positive controls, and for the coding region of the inactive MYOD1 gene and an intergeic region on chromosome 11, used as negative controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).

ChIP-Seq

ChIP was performed with 5 ug of antibody. The figure shows the peak distribution along the complete sequence and a 3 mb region of chromosome 5 and in two regions surrounding the IRS2 and ANKRD32 positive control genes. The position of the amplicon used for ChIP-qPCR is indicated by an arrow.

Gene Info — EP300		
Entrez GenelD	2033	
Protein Accession#	<u>Q09472</u>	
Gene Name	EP300	
Gene Alias	KAT3B, p300	
Gene Description	E1A binding protein p300	
Omim ID	<u>114500 180849 602700</u>	
Gene Ontology	Hyperlink	

😭 Abnova	Product Information
Gene Summary	This gene encodes the adenovirus E1A-associated cellular p300 transcriptional co-activator prot ein. It functions as histone acetyltransferase that regulates transcription via chromatin remodeling and is important in the processes of cell proliferation and differentiation. It mediates cAMP-gene r egulation by binding specifically to phosphorylated CREB protein. This gene has also been identif ied as a co-activator of HIF1A (hypoxia-inducible factor 1 alpha), and thus plays a role in the stimu lation of hypoxia-induced genes such as VEGF. Defects in this gene are a cause of Rubinstein-T aybi syndrome and may also play a role in epithelial cancer. [provided by RefSeq
Other Designations	E1A-associated protein p300 E1A-binding protein, 300kD OTTHUMP00000028668 histone acet yltransferase p300

Pathway

- Adherens junction
- <u>Cell cycle</u>
- Jak-STAT signaling pathway
- Long-term potentiation
- Melanogenesis
- Notch signaling pathway
- Pathways in cancer
- Prostate cancer
- <u>Renal cell carcinoma</u>
- TGF-beta signaling pathway
- Wnt signaling pathway

Disease

- Breast cancer
- Breast Neoplasms
- <u>Cardiovascular Diseases</u>
- Diabetes Mellitus
- Ductus Arteriosus

😵 Abnova

- Edema
- Genetic Predisposition to Disease
- Infant
- Kidney Failure
- <u>Neoplasms</u>
- Ovarian cancer
- Ovarian Neoplasms
- Spinal Dysraphism
- Thyroid Neoplasms