


MUC1 monoclonal antibody, clone GP1.4

Catalog # MAB16236 Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human breast with MUC1 monoclonal antibody, clone GP1.4 (Cat # MAB16236).

Specification

Product Description	Mouse monoclonal antibody raised against human MUC1.
Immunogen	Human milk fat globule.
Host	Mouse
Reactivity	Human
Form	Liquid
Purification	Protein G purification
Isotype	IgG1
Recommend Usage	Flow Cytometry (1:25-1:200) Immunocytochemistry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (10 ug/mL) Immunohistochemistry (Frozen sections) (1:25-1:200) Western Blot (1:100-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide).

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human breast with MUC1 monoclonal antibody, clone GP1.4 (Cat # MAB16236).
- Immunohistochemistry (Frozen sections)
- Immunocytochemistry
- Flow Cytometry

Gene Info — MUC1

Entrez GeneID [4582](#)

Protein Accession# [P15941](#)

Gene Name MUC1

Gene Alias CD227, EMA, H23AG, MAM6, PEM, PEMT, PUM

Gene Description mucin 1, cell surface associated

Omim ID [158340](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene is a member of the mucin family and encodes a membrane bound, glycosylated phosphoprotein. The protein is anchored to the apical surface of many epithelia by a transmembrane domain, with the degree of glycosylation varying with cell type. It also includes a 20 aa variable number tandem repeat (VNTR) domain, with the number of repeats varying from 20 to 120 in different individuals. The protein serves a protective function by binding to pathogens and also functions in a cell signaling capacity. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. Multiple alternatively spliced transcript variants that encode different isoforms of this gene have been reported, but the full-length nature of only some has been determined. [provided by RefSeq]

Other Designations

DF3 antigen|H23 antigen|MUC-1/SEC|MUC-1/X|MUC1/ZD|OTTHUMP00000033787|OTTHUMP00000033790|OTTHUMP00000033795|breast carcinoma-associated antigen DF3|episialin|mucin 1|mucin 1, transmembrane|peanut-reactive urinary mucin|polymorphic epithelial mucin|tumor as

Disease

- [Adenocarcinoma](#)
- [Asthma](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Carcinoma](#)
- [Chronic Disease](#)
- [Disease Susceptibility](#)
- [Gastritis](#)
- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)
- [Infant](#)
- [Kidney Failure](#)
- [Lung Diseases](#)
- [Lung Neoplasms](#)

- [Metaplasia](#)
- [NARP](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Stomach Neoplasms](#)