


PTPRC monoclonal antibody, clone E19-G

Catalog # MAB1649

Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human tonsil tissue (4 um) stained with PTPRC monoclonal antibody, clone E19-G (Cat # MAB1649) shows strong positive immunostaining. Kindly performed and provided by Katarina Poliakov, MD and Lubomir Straka, MD, Ph. D. from Clinical Pathology Presov, Ltd., Presov, Slovakia.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of PTPRC.
Immunogen	A synthetic peptide corresponding to C-terminus of human PTPRC.
Host	Rabbit
Reactivity	Human
Form	Liquid
Isotype	IgG
Quality Control Testing	Antibody Reactive Against Synthetic Peptide.
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:100-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 20 mM Tris-HCl, pH 8.0 (20 mg/mL BSA, 0.05% sodium azide)
Storage Instruction	Store at 4°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human tonsil tissue (4 um) stained with PTPRC monoclonal antibody, clone E19-G (Cat # MAB1649) shows strong positive immunostaining. Kindly performed and provided by Katarina Poliakova, MD and Lubomir Straka, MD, Ph. D. from Clinical Pathology Presov, Ltd., Presov, Slovakia.

Gene Info — PTPRC

Entrez GeneID	5788
Gene Name	PTPRC
Gene Alias	B220, CD45, CD45R, GP180, LCA, LY5, T200
Gene Description	protein tyrosine phosphatase, receptor type, C
Omim ID	126200 151460 609532
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extra cellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Four alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq]
Other Designations	CD45 antigen T200 glycoprotein T200 leukocyte common antigen glycoprotein leukocyte-common antigen protein tyrosine phosphatase, receptor type, c polypeptide

Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Primary immunodeficiency](#)

- [T cell receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Ascariasis](#)
- [Autoimmune Diseases](#)
- [Cardiomyopathy](#)
- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hashimoto Disease](#)
- [Hepatitis](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Inflammatory Bowel Diseases](#)
- [Lupus Erythematosus](#)
- [Lymphopenia](#)
- [Multiple Sclerosis](#)
- [Paraparesis](#)
- [Scleroderma](#)
- [Severe combined immunodeficiency](#)
- [Thyroiditis](#)