


RecomAb™

ITGAM monoclonal antibody, clone RM290


Catalog # MAB16920 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis of (1) HL60 cells nontreated with DMSO, (2) HL60 cell treated with DMSO.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human ITGAM.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to C-terminus of human ITGAM.
Reactivity	Human
Specificity	This antibody reacts to human CD11b (Integrin alpha-M).

Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:100-200) Western Blot (1:200-500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis of (1) HL60 cells nontreated with DMSO, (2) HL60 cell treated with DMSO.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil.

Gene Info — ITGAM

Entrez GeneID	3684
Gene Name	ITGAM
Gene Alias	CD11B, CR3A, MAC-1, MAC1A, MGC117044, MO1A, SLEB6
Gene Description	integrin, alpha M (complement component 3 receptor 3 subunit)
Omim ID	120980
Gene Ontology	Hyperlink

Gene Summary

This gene encodes the integrin alpha M chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

antigen CD11b (p170)|complement component receptor 3, alpha|integrin alpha M|macrophage antigen alpha polypeptide|neutrophil adherence receptor alpha-M subunit

Pathway

- [Cell adhesion molecules \(CAMs\)](#)
- [Hematopoietic cell lineage](#)
- [Leukocyte transendothelial migration](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Autoimmune Diseases](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)
- [Lupus Erythematosus](#)
- [Lupus Nephritis](#)
- [Macular Degeneration](#)
- [Nephritis](#)
- [Stomach Ulcer](#)