Gabbr2 monoclonal antibody, clone S81-2 (APC)


Catalog # MAB17003 Size 100 ug

Applications



Western Blot (Tissue lysate)

Western Blot analysis of rat brain membrane lysate with Gabbr2 monoclonal antibody, clone S81-2 (APC) (Cat # MAB17003).



Immunocytochemistry

Immunocytochemical staining of SK-N-BE with Gabbr2 monoclonal antibody, clone S81-2 (APC) (Cat # MAB17003). (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Gabbr2 Antibody and (D) Composite.

Specification	
Product Description	Mouse monoclonal antibody raised against partial recombinant rat Gabbr2.
Immunogen	Recombinant protein corresponding to amino acids 861-912 of rat Gabbr2.
Host	Mouse
Reactivity	Human, Rat
Form	Liquid

😵 Abnova

Product Information

Conjugation	APC
Purification	Protein G purification
lsotype	lgG1
Recommend Usage	Immunocytochemistry
	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
	Western Blot (1:1000)
	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (50% glycerol, 0.09% sodium azide).
Storage Instruction	Store at -20°C.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

• Western Blot (Tissue lysate)

Western Blot analysis of rat brain membrane lysate with Gabbr2 monoclonal antibody, clone S81-2 (APC) (Cat # MAB17003).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
- Immunocytochemistry

Immunocytochemical staining of SK-N-BE with Gabbr2 monoclonal antibody, clone S81-2 (APC) (Cat # MAB17003). (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Gabbr2 Antibody and (D) Composite.

Immunofluorescence

Gene Info — Gabbr2

Entrez GenelD	83633
Protein Accession#	<u>088871</u>
Gene Name	Gabbr2
Gene Alias	Gpr51
Gene Description	gamma-aminobutyric acid (GABA) B receptor 2



Product Information

Gene Ontology	Hyperlink
Other Designations	G protein-coupled receptor 51 GABA-B R2 receptor

Publication Reference

• A single subunit (GB2) is required for G-protein activation by the heterodimeric GABA(B) receptor.

Duthey B, Caudron S, Perroy J, Bettler B, Fagni L, Pin JP, Prezeau L. The Journal of Biological Chemistry 2002 Feb; 277(5):236.