


BCL2L10 monoclonal antibody, clone 8A11G12


Catalog # MAB17524 Size 100 ug

Applications




Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) BCL2L10-hlgGfc transfected HEK293 cell lysate.



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of BCL2L10 monoclonal antibody, clone 8A11G12.



Flow Cytometry

Flow cytometric analysis of Hela cells with BCL2L10 monoclonal antibody (green) and negative control (red).

Specification

Product Description

Mouse monoclonal antibody raised against recombinant human BCL2L10.

Immunogen	Recombinant protein corresponding to amino acid 31-186 of human BCL2L10 from <i>E. coli</i> .
Host	Mouse
Theoretical MW (kDa)	22
Reactivity	Human
Form	Liquid
Isotype	IgG2a
Recommend Usage	ELISA (1:10000) Western Blot (1:500-1:2000) Immunocytochemistry Flow Cytometry (1:200-1:400) Immunohistochemistry The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) BCL2L10-hlgGFc transfected HEK293 cell lysate.

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis of BCL2L10 monoclonal antibody, clone 8A11G12.

- Flow Cytometry

Flow cytometric analysis of Hela cells with BCL2L10 monoclonal antibody (green) and negative control (red).

Gene Info — BCL2L10

Entrez GeneID	10017
Gene Name	BCL2L10
Gene Alias	BCL-B, Boo, Diva, MGC129810, MGC129811

Gene Description	BCL2-like 10 (apoptosis facilitator)
Omim ID	606910
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains conserved BH4, BH1 and BH2 domains. This protein can interact with other members of BCL-2 protein family including BCL2, BCL2L1/BCL-X(L), and BAX. Overexpression of this gene has been shown to suppress cell apoptosis possibly through the prevention of cytochrome C release from the mitochondria, and thus activating caspase-3 activation. The mouse counterpart of this protein is found to interact with Apaf1 and forms a protein complex with Caspase 9, which suggests the involvement of this protein in APAF1 and CASPASE 9 related apoptotic pathway. [provided by RefSeq]
Other Designations	-

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Occupational Diseases](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)