


DNM2 monoclonal antibody, clone 3F5C7


Catalog # MAB17595 Size 100 ug

Applications




Western Blot (Cell lysate)

Western blot analysis of (1) U251 cell, (2) Hela cell, (3) K562 cell with DNM2 monoclonal antibody.



Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) DNM2-hlgGFc transfected HEK293 cell lysate.



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of DNM2 monoclonal antibody, clone 3F5C7.

Specification

Product Description

Mouse monoclonal antibody raised against recombinant human DNM2.



Product Information

Immunogen	Recombinant protein corresponding to amino acid 520-744 of human DNM2 from E. coli.
Host	Mouse
Theoretical MW (kDa)	98
Reactivity	Human
Form	Liquid
Isotype	lgG2a
Recommend Usage	ELISA (1:10000) Western Blot (1:500-1:2000) Immunocytochemistry Flow Cytometry Immunohistochemistry The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western blot analysis of (1) U251 cell, (2) Hela cell, (3) K562 cell with DNM2 monoclonal antibody.

Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) DNM2-hlgGFc transfected HEK293 cell lysate.

Enzyme-linked Immunoabsorbent Assay

ELISA analysis of DNM2 monoclonal antibody, clone 3F5C7.

Gene	6 _	
I-ODO	IDTO	1 11/11/11/11

Entrez GeneID	<u>1785</u>
Gene Name	DNM2
Gene Alias	CMTDI1, CMTDIB, DI-CMTB, DYN2, DYNII



Product Information

Gene Description	dynamin 2
Omim ID	<u>160150</u> <u>602378</u> <u>606482</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Dynamins represent one of the subfamilies of GTP-binding proteins. These proteins share considerable sequence similarity over the N-terminal portion of the molecule, which contains the GTP as e domain. Dynamins are associated with microtubules. They have been implicated in cell process es such as endocytosis and cell motility, and in alterations of the membrane that accompany certa in activities such as bone resorption by osteoclasts. Dynamins bind many proteins that bind actinand other cytoskeletal proteins. Dynamins can also self-assemble, a process that stimulates GTP ase activity. Four alternatively spliced transcripts encoding different proteins have been described. Additional alternatively spliced transcripts may exist, but their full-length nature has not been determined. [provided by RefSeq
Other Designations	dynamin II

Pathway

- Endocytosis
- Fc gamma R-mediated phagocytosis

Disease

- Alzheimer disease
- Anorexia Nervosa
- Bulimia
- Genetic Predisposition to Disease