


CXCR3 monoclonal antibody, clone 5C10E6

Catalog # MAB17606


Size 100 ug

Applications




Western Blot (Cell lysate)

Western blot analysis of (1) Hela cell, (2) L-02 cell with CXCR3 monoclonal antibody.




Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) CXCR3-hlgGfc transfected HEK293 cell lysate.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded esophageal cancer tissues with CXCR3 monoclonal antibody.



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of CXCR3 monoclonal antibody, clone 5C10E6.



Flow Cytometry

Flow cytometric analysis of Jurkat cells with CXCR3 monoclonal antibody (green) and negative control (red).

Specification

Product Description	Mouse monoclonal antibody raised against recombinant human CXCR3.
Immunogen	Recombinant protein corresponding to amino acid of human CXCR3 from <i>E. coli</i> .
Host	Mouse
Theoretical MW (kDa)	40.7
Reactivity	Human
Form	Liquid
Isotype	IgG1
Recommend Usage	ELISA (1:10000) Western Blot (1:500-1:2000) Immunocytochemistry Flow Cytometry (1:200-1:400) Immunohistochemistry (1:200-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.05% sodium azide).

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of (1) Hela cell, (2) L-02 cell with CXCR3 monoclonal antibody.

- Western Blot (Transfected lysate)

Western blot analysis of (1) HEK293 cells, (2) CXCR3-hlgGfc transfected HEK293 cell lysate.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded esophageal cancer tissues with CXCR3 monoclonal antibody.

- Enzyme-linked Immunoabsorbent Assay

ELISA analysis of CXCR3 monoclonal antibody, clone 5C10E6.

- Flow Cytometry

Flow cytometric analysis of Jurkat cells with CXCR3 monoclonal antibody (green) and negative control (red).

Gene Info — CXCR3

Entrez GeneID[2833](#)**Gene Name**

CXCR3

Gene Alias

CD182, CD183, CKR-L2, CMKAR3, GPR9, IP10-R, Mig-R, MigR

Gene Description

chemokine (C-X-C motif) receptor 3

Omim ID[300574](#)**Gene Ontology**[Hyperlink](#)

Gene Summary

This gene encodes a G protein-coupled receptor with selectivity for three chemokines, termed IP10 (interferon-g-inducible 10 kDa protein), Mig (monokine induced by interferon-g) and I-TAC (interferon-inducible T cell a-chemoattractant). IP10, Mig and I-TAC belong to the structural subfamily of CXC chemokines, in which a single amino acid residue separates the first two of four highly conserved Cys residues. Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic, most notably integrin activation, cytoskeletal changes and chemotactic migration. Inhibition by Bordetella pertussis toxin suggests that heterotrimeric G protein of the Gi-subclass couple to this protein. Signal transduction has not been further analyzed but may include the same enzymes that were identified in the signaling cascade induced by other chemokine receptors. As a consequence of chemokine-induced cellular desensitization (phosphorylation-dependent receptor internalization), cellular responses are typically rapid and short in duration. Cellular responsiveness is restored after dephosphorylation of intracellular receptors and subsequent recycling to the cell surface. This gene is prominently expressed in in vitro cultured effector/memory T cells, and in T cells present in many types of inflamed tissues. In addition, IP10, Mig and I-TAC are commonly produced by local cells in inflammatory lesion, suggesting that this gene and its chemokines participate in the recruitment of inflammatory cells. Therefore, this protein is a target for the development of small molecular weight antagonists, which may be used in the treatment of diverse inflammatory diseases. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

G protein-coupled receptor 9|IP10 receptor|Mig receptor|OTTHUMP00000070257|chemokine (C-X-C) receptor 3

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)

Disease

- [Asthma](#)
- [Bronchiolitis](#)
- [Coronary Artery Disease](#)
- [Genetic Predisposition to Disease](#)
- [Infant](#)
- [Respiratory Syncytial Virus Infections](#)