


BAD (phospho S112) monoclonal antibody, clone B9

Catalog # MAB18806 Size 20 uL

Applications



Flow Cytometry

Flow cytometric analysis of U937 cells with BAD (phospho S112) monoclonal antibody, clone B9 (Cat # MAB18806). Secondary antibody only negative control (blue) or untreated (red) or treated with Calyculin A (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human BAD.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding S112 of human BAD.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Protein A/G purification
Isotype	IgG1, lamda
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (50% glycerol, 0.1% BSA and 0.02% sodium azide).
Storage Instruction	Store at -20°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of U937 cells with BAD (phospho S112) monoclonal antibody, clone B9 (Cat # MAB18806). Secondary antibody only negative control (blue) or untreated (red) or treated with Calyculin A (green).

Gene Info — BAD

Entrez GeneID	572
---------------	---------------------

Gene Name	BAD
-----------	-----

Gene Alias	BBC2, BCL2L8
------------	--------------

Gene Description	BCL2-associated agonist of cell death
------------------	---------------------------------------

Omim ID	603167
---------	------------------------

Gene Ontology	Hyperlink
---------------	---------------------------

Gene Summary	The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq]
--------------	--

Other Designations	BCL-X/BCL-2 binding protein BCL2-antagonist of cell death protein BCL2-binding component 6 BCL2-binding protein
--------------------	---

Pathway

- [Acute myeloid leukemia](#)
- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Chronic myeloid leukemia](#)

- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Insulin signaling pathway](#)
- [Melanoma](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [VEGF signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lymphoma](#)
- [Parkinson disease](#)
- [Thyroid Neoplasms](#)