
## CBL (phospho Y700) monoclonal antibody, clone E1 (APC)

Catalog # MAB18837 Size 100 Reactions

## Applications



## Flow Cytometry

Flow cytometric analysis of C6 cells with CBL (phospho Y700) monoclonal antibody, clone E1 (APC) (Cat # MAB18837). Cell only negative control (blue) or treated with imatinib (red) or with pervanadate (green).

Specification	
Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human CBL.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y700 of human CBL.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Form	Liquid
Conjugation	APC
Purification	Protein A/G purification
lsotype	lgG1, kappa
Recommend Usage	Flow Cytometry (5 uL/10 <sup>6</sup> cells) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).
Storage Instruction	Store at 4°C.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😵 Abnova

## **Product Information**

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## Applications

#### Flow Cytometry

Flow cytometric analysis of C6 cells with CBL (phospho Y700) monoclonal antibody, clone E1 (APC) (Cat # MAB18837). Cell only negative control (blue) or treated with imatinib (red) or with pervanadate (green).

Gene Info — CBL	
Entrez GenelD	<u>867</u>
Gene Name	CBL
Gene Alias	C-CBL, CBL2, RNF55
Gene Description	Cas-Br-M (murine) ecotropic retroviral transforming sequence
Omim ID	<u>165360</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The cbl oncogene was first identified as part of a transforming retrovirus which induces mouse pr e-B and pro-B cell lymphomas. As an adaptor protein for receptor protein-tyrosine kinases, it posi tively regulates receptor protein-tyrosine kinase ubiquitination in a manner dependent upon its var iant SH2 and RING finger domains. Ubiquitination of receptor protein-tyrosine kinases terminates signaling by marking active receptors for degradation. [provided by RefSeq
Other Designations	oncogene CBL2

## Pathway

- <u>Chronic myeloid leukemia</u>
- Endocytosis
- ErbB signaling pathway
- Insulin signaling pathway
- Jak-STAT signaling pathway
- Pathways in cancer

# 😵 Abnova

## **Product Information**

- <u>T cell receptor signaling pathway</u>
- Ubiquitin mediated proteolysis

#### Disease

- Diabetes Mellitus
- Disease Progression
- Genetic Predisposition to Disease
- Leukemia
- Tobacco Use Disorder