


# CBL (phospho Y700) monoclonal antibody, clone E1 (PE)

Catalog # MAB18840      Size 10 Reactions

## Applications



### Flow Cytometry

Flow cytometric analysis of C6 cells with CBL (phospho Y700) monoclonal antibody, clone E1 (PE) (Cat # MAB18840). Cell only negative control (blue) or treated with imatinib (red) or with pervanadate (green).

## Specification

|                            |  |
|----------------------------|--|
| <b>Product Description</b> | Rabbit monoclonal antibody raised against synthetic phosphopeptide of human CBL.                               |
| <b>Immunogen</b>           | A synthetic phosphopeptide corresponding to residues surrounding Y700 of human CBL.                            |
| <b>Host</b>                | Rabbit   |
| <b>Reactivity</b>          | Human, Mouse, Rat  |
| <b>Form</b>                | Liquid   |
| <b>Conjugation</b>         | PE   |
| <b>Purification</b>        | Protein A/G purification   |
| <b>Isotype</b>             | IgG1, kappa  |
| <b>Recommend Usage</b>     | Flow Cytometry (5 $\mu$ L/ $10^6$ cells)<br>The optimal working dilution should be determined by the end user. |
| <b>Storage Buffer</b>      | In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).   |

**Storage Instruction**

Store at 4°C.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Flow Cytometry

Flow cytometric analysis of C6 cells with CBL (phospho Y700) monoclonal antibody, clone E1 (PE) (Cat # MAB18840). Cell only negative control (blue) or treated with imatinib (red) or with pervanadate (green).

## Gene Info — CBL

**Entrez GeneID**[867](#)**Gene Name**

CBL

**Gene Alias**

C-CBL, CBL2, RNF55

**Gene Description**

Cas-Br-M (murine) ecotropic retroviral transforming sequence

**Omim ID**[165360](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The cbl oncogene was first identified as part of a transforming retrovirus which induces mouse pre-B and pro-B cell lymphomas. As an adaptor protein for receptor protein-tyrosine kinases, it positively regulates receptor protein-tyrosine kinase ubiquitination in a manner dependent upon its variant SH2 and RING finger domains. Ubiquitination of receptor protein-tyrosine kinases terminates signaling by marking active receptors for degradation. [provided by RefSeq]

**Other Designations**

oncogene CBL2

## Pathway

- [Chronic myeloid leukemia](#)
- [Endocytosis](#)
- [ErbB signaling pathway](#)
- [Insulin signaling pathway](#)
- [Jak-STAT signaling pathway](#)

- [Pathways in cancer](#)
- [T cell receptor signaling pathway](#)
- [Ubiquitin mediated proteolysis](#)

## Disease

- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Genetic Predisposition to Disease](#)
- [Leukemia](#)
- [Tobacco Use Disorder](#)