


CBL (phospho Y774) monoclonal antibody, clone R3B8 (PE)

Catalog # MAB18847 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of Daudi cells with CBL (phospho Y774) monoclonal antibody, clone R3B8 (PE) (Cat # MAB18847). Cell only (blue) or untreated (red) or treated with IFN α + IL-4 + pervanadate (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human CBL.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y774 of human CBL.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Conjugation	PE
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (5 μ L/ 10^6 cells) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).
Storage Instruction	Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of Daudi cells with CBL (phospho Y774) monoclonal antibody, clone R3B8 (PE) (Cat # MAB18847). Cell only (blue) or untreated (red) or treated with IFN α + IL-4 + pervanadate (green).

Gene Info — CBL

Entrez GeneID	867
---------------	---------------------

Gene Name	CBL
-----------	-----

Gene Alias	C-CBL, CBL2, RNF55
------------	--------------------

Gene Description	Cas-Br-M (murine) ecotropic retroviral transforming sequence
------------------	--

Omim ID	165360
---------	------------------------

Gene Ontology	Hyperlink
---------------	---------------------------

Gene Summary	The cbl oncogene was first identified as part of a transforming retrovirus which induces mouse pre-B and pro-B cell lymphomas. As an adaptor protein for receptor protein-tyrosine kinases, it positively regulates receptor protein-tyrosine kinase ubiquitination in a manner dependent upon its variant SH2 and RING finger domains. Ubiquitination of receptor protein-tyrosine kinases terminates signaling by marking active receptors for degradation. [provided by RefSeq]
--------------	--

Other Designations	oncogene CBL2
--------------------	---------------

Pathway

- [Chronic myeloid leukemia](#)
- [Endocytosis](#)
- [ErbB signaling pathway](#)
- [Insulin signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pathways in cancer](#)

- [T cell receptor signaling pathway](#)
- [Ubiquitin mediated proteolysis](#)

Disease

- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Genetic Predisposition to Disease](#)
- [Leukemia](#)
- [Tobacco Use Disorder](#)