


STAT1 (phospho Y701) monoclonal antibody, clone 3E6 (FITC)

Catalog # MAB19120 Size 10 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of U937 cells with STAT1 (phospho Y701) monoclonal antibody, clone 3E6 (FITC) (Cat # MAB19120). Unstained untreated cells as negative control (blue) or stained untreated (red) or treated with IFN α , IL-4 and pervanadate (green).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human STAT1.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y701 of human STAT1.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Conjugation	FITC
Purification	Protein A/G purification
Isotype	IgG1, kappa
Recommend Usage	Flow Cytometry (5 μ L/ 10^6 cells) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (0.2% BSA, 0.09% sodium azide).

Storage Instruction

Store at 4°C.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of U937 cells with STAT1 (phospho Y701) monoclonal antibody, clone 3E6 (FITC) (Cat # MAB19120). Unstained untreated cells as negative control (blue) or stained untreated (red) or treated with IFN α , IL-4 and pervanadate (green).

Gene Info — STAT1

Entrez GeneID[6772](#)**Gene Name**

STAT1

Gene Alias

DKFZp686B04100, ISGF-3, STAT91

Gene Description

signal transducer and activator of transcription 1, 91kDa

Omim ID[209950](#) [600555](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations

OTTHUMP00000165047|signal transducer and activator of transcription 1|signal transducer and activator of transcription-1|transcription factor ISGF-3 components p91/p84

Pathway

- [Chemokine signaling pathway](#)
- [Jak-STAT signaling pathway](#)

- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Campylobacter Infections](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Glomerulonephritis](#)
- [Hepatitis B](#)
- [Hepatitis C](#)

- [Hypersensitivity](#)
- [Infant](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neutropenia](#)
- [Osteoporosis](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Proteinuria](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Salmonella Infections](#)
- [Thrombocytopenia](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)

- [Werner syndrome](#)