


BCL2 monoclonal antibody, clone HF-2


Catalog # MAB19606 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of (1) Jurkat cell lysate;(2) MCF-7 cell lysate with BCL2 monoclonal antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human spleen with BCL2 monoclonal antibody.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human BCL2.
Immunogen	A synthetic peptide corresponding to human BCL2.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Purification	Affinity purification
Isotype	IgG

Recommend Usage	Immunocytochemistry (1:50-1:100) Immunofluorescence (1:50-1:100) Immunohistochemistry (1:50-1:100) Flow Cytometry (1:50) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of (1) Jurkat cell lysate;(2) MCF-7 cell lysate with BCL2 monoclonal antibody.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human spleen with BCL2 monoclonal antibody.

- Immunocytochemistry

- Immunofluorescence

- Immunoprecipitation

- Flow Cytometry

Gene Info — BCL2

Entrez GeneID	596
Protein Accession#	P10415
Gene Name	BCL2
Gene Alias	Bcl-2

Gene Description	B-cell CLL/lymphoma 2
Omim ID	151430
Gene Ontology	Hyperlink
Gene Summary	This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends. [provided by RefSeq]
Other Designations	B-cell lymphoma protein 2 OTTHUMP00000163680

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Focal adhesion](#)
- [Neurotrophin signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer Disease](#)
- [Amnesia](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Azoospermia](#)
- [Bile Duct Neoplasms](#)

- [Brain Injuries](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cholangiocarcinoma](#)
- [Chromosome Aberrations](#)
- [Clubfoot](#)
- [Cognition](#)
- [Cognition Disorders](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Kidney Neoplasms](#)
- [Laryngeal Neoplasms](#)
- [Leukemia](#)
- [Leukemia-Lymphoma](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)

- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [NARP](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Recurrence](#)
- [Neovascularization](#)
- [Neuropsychological Tests](#)
- [Occupational Diseases](#)
- [Oropharyngeal Neoplasms](#)
- [Ovarian Neoplasms](#)
- [Postoperative Complications](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Prosthesis Failure](#)
- [Pulmonary Disease](#)
- [Thyroid Neoplasms](#)
- [Translocation](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)