


CDC2 monoclonal antibody, clone AAD-3

Catalog # MAB19873 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of HeLa cell lysate with CDC2 monoclonal antibody.

| Specification | |
|---------------------|--|
| Product Description | Rabbit monoclonal antibody raised against synthetic peptide of human CDC2. |
| Immunogen | A synthetic peptide corresponding to human CDC2. |
| Host | Rabbit |
| Reactivity | Human |
| Form | Liquid |
| Purification | Affinity purification |
| Isotype | lgG |
| Recommend Usage | Immunoprecipitation (1:50) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In PBS, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage Instruction | Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and st ored frozen at -20°C for a longer time. Avoid repeated freezing and thawing. |



Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western blot analysis of HeLa cell lysate with CDC2 monoclonal antibody.

Immunoprecipitation

| Gene Info — CDC2 | |
|--------------------|--|
| Entrez GenelD | 983 |
| Protein Accession# | P06493 |
| Gene Name | CDC2 |
| Gene Alias | CDC28A, CDK1, DKFZp686L20222, MGC111195 |
| Gene Description | cell division cycle 2, G1 to S and G2 to M |
| Omim ID | <u>116940</u> |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting f actor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitot ic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phos phorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq |
| Other Designations | OTTHUMP00000019660 cell cycle controller CDC2 cell division control protein 2 homolog cell division cycle 2 protein cyclin-dependent kinase 1 p34 protein kinase |

Pathway

- Cell cycle
- Gap junction



p53 signaling pathway

Disease

- Alzheimer disease
- Breast cancer
- Breast Neoplasms
- Dementia
- Genetic Predisposition to Disease
- Lung Neoplasms
- Pulmonary Disease