


# Histone H2A (hydroxyl Y39) monoclonal antibody, clone DED-8

Catalog # MAB20142

Size 100 uL

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of mouse colon with Histone H2A (hydroxyl Y39) monoclonal antibody, clone DED-8 (Cat # MAB20142).

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against synthetic hydroxyl peptide of Histone H2A (hydroxyl Y39).
<b>Immunogen</b>	A synthetic hydroxyl peptide corresponding to residues surrounding Y39 of Histone H2A.
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Purification</b>	Affinity purification
<b>Isotype</b>	IgG
<b>Recommend Usage</b>	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:500-1:2000) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).
<b>Storage Instruction</b>	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

## Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of mouse colon with Histone H2A (hydroxyl Y39) monoclonal antibody, clone DED-8 (Cat # MAB20142).