


HLA-DRA monoclonal antibody, clone FGI-8

Catalog # MAB20167 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis of Raji cell lysate.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human HLA-DRA.
Immunogen	A synthetic peptide corresponding to human HLA-DRA.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Flow Cytometry (1:50) Immunocytochemistry (1:50-200) Immunofluorescence (1:50-200) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-200) Western Blot (1:1000-5000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot (Cell lysate) analysis of Raji cell lysate.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

- Immunocytochemistry

- Immunofluorescence

- Flow Cytometry

Gene Info — HLA-DRA

Entrez GeneID[3122](#)**Protein Accession#**[P01903](#)**Gene Name**

HLA-DRA

Gene Alias

HLA-DRA1

Gene Description

major histocompatibility complex, class II, DR alpha

Omim ID[142860](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

HLA-DRA is one of the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha and a beta chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. DRA does not have polymorphisms in the peptide binding part and acts as the sole alpha chain for DRB1, DRB3, DRB4 and DRB5. [provided by RefSeq]

Other Designations

HLA class II histocompatibility antigen, DR alpha chain|MHC cell surface glycoprotein|histocompatibility antigen HLA-DR alpha

Pathway

- [Allograft rejection](#)
- [Antigen processing and presentation](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Hematopoietic cell lineage](#)
- [Systemic lupus erythematosus](#)
- [Type I diabetes mellitus](#)

Disease

- [Abortion](#)
- [Alzheimer disease](#)
- [Anticipation](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Atrial Fibrillation](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)

- [Cerebral Amyloid Angiopathy](#)
- [Cerebrovascular Accident](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Drug Hypersensitivity](#)
- [Edema](#)
- [Enterovirus Infections](#)
- [Food Hypersensitivity](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Hypersensitivity](#)
- [IgA Deficiency](#)
- [Latex Hypersensitivity](#)
- [Lichen Planus](#)
- [Lupus Erythematosus](#)
- [Multiple Sclerosis](#)
- [Myasthenia Gravis](#)
- [Neuroblastoma](#)
- [Obesity](#)
- [Parkinson Disease](#)
- [Respiratory Tract Infections](#)
- [Sarcoidosis](#)

- [Spondylitis](#)
- [Uveomeningoencephalitic Syndrome](#)