
# TSPO monoclonal antibody, clone DAG-20


Catalog # MAB20299 Size 100 uL

## Applications



#### Western Blot (Cell lysate)

Western blot analysis of (1) U87-MG cell lysate; (2) A431 cell lysate with TSPO monoclonal antibody.



### Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining of paraffin-embedded human colon with TSPO monoclonal antibody.

Specification	
Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human TSPO.
Immunogen	A synthetic peptide corresponding to human TSPO.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Affinity purification
lsotype	lgG



Recommend Usage	Flow Cytometry (1:50) Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Immunohistochemistry (1:50-1:200) Immunoprecipitation (1:50) Western Blot (1:5000-1:10000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and st ored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## Applications

• Western Blot (Cell lysate)

Western blot analysis of (1) U87-MG cell lysate; (2) A431 cell lysate with TSPO monoclonal antibody.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human colon with TSPO monoclonal antibody.

- Immunocytochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytometry

Gene Info — TSPO		
Entrez GenelD	<u>706</u>	
Protein Accession#	<u>P30536</u>	
Gene Name	TSPO	
Gene Alias	BZRP, DBI, IBP, MBR, PBR, PKBS, PTBR, mDRC, pk18	

😵 Abnova

**Product Information** 

Gene Description	translocator protein (18kDa)
Omim ID	<u>109610</u>
Gene Ontology	Hyperlink
Gene Summary	Present mainly in the mitochondrial compartment of peripheral tissues, the protein encoded by thi s gene interacts with some benzodiazepines and has different affinities than its endogenous coun terpart. The protein is a key factor in the flow of cholesterol into mitochondria to permit the initiatio n of steroid hormone synthesis. Alternate transcriptional splice variants, encoding different isofor ms, have been characterized. [provided by RefSeq
Other Designations	OTTHUMP00000028891 benzodiazepine peripheral binding site mitochondrial benzodiazepine r eceptor peripheral benzodiazepine receptor peripheral-type benzodiazepine receptor translocator protein

## Pathway

<u>Neuroactive ligand-receptor interaction</u>

#### Disease

- Asperger Syndrome
- <u>Autistic Disorder</u>
- Genetic Predisposition to Disease
- <u>Kidney Failure</u>
- Lung Neoplasms
- Mood Disorders
- Panic Disorder
- Social Perception
- Urinary Bladder Neoplasms