


SNCA (phospho S129) monoclonal antibody, clone BBF-19

Catalog # MAB20324 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of (1) 293T cell lysate; (2) 293T cell lysate transfected with Polo-Like Kinase 2 and alpha Synuclein with SNCA (phospho S129) monoclonal antibody.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human SNCA.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding S129 of human SNCA.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Immunohistochemistry (1:50-1:200) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.4-0.5 mg/mL BSA, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of (1) 293T cell lysate; (2) 293T cell lysate transfected with Polo-Like Kinase 2 and alpha Synuclein with SNCA (phospho S129) monoclonal antibody.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

- Immunocytochemistry

- Immunofluorescence

Gene Info — SNCA

Entrez GeneID[6622](#)**Protein Accession#**[P37840](#)**Gene Name**

SNCA

Gene Alias

MGC110988, NACP, PARK1, PARK4, PD1

Gene Description

synuclein, alpha (non A4 component of amyloid precursor)

Omim ID[127750](#) [163890](#) [168601](#) [605543](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Four alternatively spliced transcripts encoding two different isoforms have been identified for this gene. [provided by RefSeq]

Other Designations

OTTHUMP00000161559|OTTHUMP00000161561|alpha synuclein|alpha-synuclein|alpha-synuclein, isoform NACP140|non A-beta component of AD amyloid|non A4 component of amyloid

Disease

- [Alcoholism](#)
- [Alzheimer disease](#)
- [Amphetamine-Related Disorders](#)
- [Association Learning](#)
- [Attention](#)
- [Cardiovascular Diseases](#)
- [Chromosome Breakage](#)
- [Cognition Disorders](#)
- [Color Perception](#)
- [Cystic fibrosis](#)
- [Dementia](#)
- [Diabetes Complications](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Essential tremor](#)
- [Genetic Predisposition to Disease](#)
- [Lewy Body Disease](#)
- [Mental Recall](#)
- [Metabolic Syndrome X](#)
- [Movement Disorders](#)
- [Multiple System Atrophy](#)
- [Neoplasms](#)
- [Nerve Degeneration](#)
- [Neuroblastoma](#)

- [Neurodegenerative Diseases](#)
- [Neuropsychological Tests](#)
- [Olfaction Disorders](#)
- [Osteoporosis](#)
- [Parkinson disease](#)
- [Parkinsonian Disorders](#)
- [Psychological Tests](#)
- [Psychomotor Performance](#)
- [Psychoses](#)
- [Serial Learning](#)
- [Smell](#)
- [Substance-Related Disorders](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)