


IKBKB monoclonal antibody, clone FOI-9

Catalog # MAB20369

Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of Daudi cell lysate with IKBKB monoclonal antibody, clone FOI-9 (Cat # MAB20369).

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human IKBKB.

Immunogen A synthetic peptide corresponding to human IKBKB.

Host Rabbit

Theoretical MW (kDa) 86.564

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage
Immunoprecipitation (1:50)
Western Blot (1:500-1:2000)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of Daudi cell lysate with IKBKB monoclonal antibody, clone FOI-9 (Cat # MAB20369).

- Immunoprecipitation

Gene Info — IKBKB

Entrez GeneID[3551](#)**Protein Accession#**[O14920](#)**Gene Name**

IKBKB

Gene Alias

FLJ40509, IKK-beta, IKK2, IKKB, MGC131801, NFKBIKB

Gene Description

inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta

Omim ID[603258](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM 164014), or RELB (MIM 604758) to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]

Other Designations

inhibitor of nuclear factor kappa B kinase beta subunit|nuclear factor NF-kappa-B inhibitor kinase beta

Pathway

- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)
- [Apoptosis](#)
- [B cell receptor signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Chronic myeloid leukemia](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [Insulin signaling pathway](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Toll-like receptor signaling pathway](#)
- [Type II diabetes mellitus](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Bronchiolitis](#)
- [Colonic Neoplasms](#)
- [Disease Susceptibility](#)

- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Inflammation](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Rectal Neoplasms](#)
- [Respiratory Syncytial Virus Infections](#)
- [Thyroid Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)