


MET monoclonal antibody, clone BFA-13


Catalog # MAB20486 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of 293 cell lysate with MET monoclonal antibody, clone BFA-13 (Cat # MAB20486).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human stomach with MET monoclonal antibody, clone BFA-13 (Cat # MAB20486).

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic peptide of human MET.
Immunogen	A synthetic peptide corresponding to human MET.
Host	Rabbit
Theoretical MW (kDa)	155.541
Reactivity	Human
Form	Liquid
Purification	Affinity purification

Isotype	IgG
Recommend Usage	Flow Cytometry (1:50) Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-1:200) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of 293 cell lysate with MET monoclonal antibody, clone BFA-13 (Cat # MAB20486).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human stomach with MET monoclonal antibody, clone BFA-13 (Cat # MAB20486).

- Immunocytochemistry

- Immunofluorescence

- Flow Cytometry

Gene Info — MET

Entrez GeneID	4233
Protein Accession#	P08581
Gene Name	MET
Gene Alias	AUTS9, HGFR, RCCP2, c-Met
Gene Description	met proto-oncogene (hepatocyte growth factor receptor)

Omim ID [114550 164860 605074 611015](#)

Gene Ontology [Hyperlink](#)

Gene Summary The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations HGF receptor|OTTHUMP00000069168|SF receptor|met proto-oncogene|met proto-oncogene tyrosine kinase|oncogene MET|scatter factor receptor

Pathway

- [Adherens junction](#)
- [Axon guidance](#)
- [Colorectal cancer](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [Focal adhesion](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Renal cell carcinoma](#)

Disease

- [Adenocarcinoma](#)
- [Autistic Disorder](#)
- [Carcinoma](#)
- [Cell Transformation](#)
- [Child Development Disorders](#)

- [Chronic Disease](#)
- [Cognition Disorders](#)
- [Disease Progression](#)
- [Gastrointestinal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hyperparathyroidism](#)
- [Lung Neoplasms](#)
- [Myopia](#)
- [Nasal Polyps](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Neuropsychological Tests](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Refractive Errors](#)
- [Rhinitis](#)
- [Schizophrenia](#)
- [Sinusitis](#)
- [Skin Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)