


# PTK2 (phospho Y397) monoclonal antibody, clone EFO-16

Catalog # MAB20515

Size 100 uL

## Applications



### Western Blot (Cell lysate)

Western Blot analysis of EGF treated 293 whole cell lysates, using PTK2 (phospho Y397) monoclonal antibody, clone EFO-16.

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human PTK2.
<b>Immunogen</b>	A synthetic phosphopeptide corresponding to residues surrounding Y397 of human PTK2.
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Purification</b>	Affinity purification
<b>Isotype</b>	IgG
<b>Recommend Usage</b>	Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.4-0.5 mg/mL BSA, 0.02% sodium azide).

**Storage Instruction**

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Note**

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot (Cell lysate)

Western Blot analysis of EGF treated 293 whole cell lysates, using PTK2 (phospho Y397) monoclonal antibody, clone EFO-16.

- Immunocytochemistry

- Immunofluorescence

## Gene Info — PTK2

**Entrez GeneID**[5747](#)**Protein Accession#**[Q05397](#)**Gene Name**

PTK2

**Gene Alias**

FADK, FAK, FAK1, pp125FAK

**Gene Description**

PTK2 protein tyrosine kinase 2

**Omim ID**[600758](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. At least four transcript variants encoding four different isoforms have been found for this gene, but the full-length nature of only two of them have been determined. [provided by RefSeq]

**Other Designations**

focal adhesion kinase 1

## Pathway

- [Axon guidance](#)
- [Chemokine signaling pathway](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Leukocyte transendothelial migration](#)
- [Pathways in cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Small cell lung cancer](#)
- [VEGF signaling pathway](#)

## Disease

- [Autistic Disorder](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)
- [Leukemia](#)
- [Mental Retardation](#)
- [Neovascularization](#)
- [Psychotic Disorders](#)
- [Schizophrenia](#)