


NBN (phospho S343) monoclonal antibody, clone BGI-14

Catalog # MAB20547 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of Jurkat cell lysate treated with Etoposide using NBN (phospho S343) monoclonal antibody, clone BGI-14.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human NBN.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding S343 of human NBN.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Immunoprecipitation (1:50) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.4-0.5 mg/mL BSA, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of Jurkat cell lysate treated with Etoposide using NBN (phospho S343) monoclonal antibody, clone BGI-14.

- Immunocytochemistry

- Immunofluorescence

- Immunoprecipitation

Gene Info — NBN

Entrez GeneID[4683](#)**Protein Accession#**[O60934](#)**Gene Name**

NBN

Gene Alias

AT-V1, AT-V2, ATV, FLJ10155, MGC87362, NBS, NBS1, P95

Gene Description

nibrin

Omim ID[251260](#) [602667](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Mutations in this gene are associated with Nijmegen breakage syndrome, an autosomal recessive chromosomal instability syndrome characterized by microcephaly, growth retardation, immunodeficiency, and cancer predisposition. The encoded protein is a member of the MRE11/RAD50 double-strand break repair complex which consists of 5 proteins. This gene product is thought to be involved in DNA double-strand break repair and DNA damage-induced checkpoint activation. [provided by RefSeq]

Other Designations

Nijmegen breakage syndrome 1 (nibrin)|cell cycle regulatory protein p95|p95 protein of the MRE11/RAD50 complex

Pathway

- [Homologous recombination](#)

Disease

- [Acute Disease](#)
- [Adenocarcinoma](#)
- [Astrocytoma](#)
- [Ataxia telangiectasia](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Central Nervous System Neoplasms](#)
- [Chromosomal Instability](#)
- [Chromosome Aberrations](#)
- [Chromosome Breakage](#)
- [Chromosome Deletion](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [DNA Damage](#)
- [DNA Repair-Deficiency Disorders](#)
- [Esophageal Neoplasms](#)

- [Gastrointestinal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genital Neoplasms](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hematologic Diseases](#)
- [Hematologic Neoplasms](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Laryngeal Neoplasms](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Malignant melanoma](#)
- [Medulloblastoma](#)
- [Melanoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mouth Neoplasms](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Neuroma](#)
- [Nijmegen Breakage Syndrome](#)
- [Obesity](#)
- [Occupational Diseases](#)

- [Ovarian cancer](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Pancreatic Neoplasms](#)
- [Parkinson disease](#)
- [Polycystic Ovary Syndrome](#)
- [Precancerous Conditions](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Puberty](#)
- [Pulmonary Disease](#)
- [Skin Neoplasms](#)
- [Stomach Neoplasms](#)
- [Syndrome](#)
- [Thrombophilia](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)