


STAT1 (phospho Y701) monoclonal antibody, clone HHB-19

Catalog # MAB20577 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of (1) A431 cell lysate, (2) A431 lysate cell treated with EGF using STAT1 (phospho Y701) monoclonal antibody, clone HHB-19.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human STAT1.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Y701 of human STAT1.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Western Blot (1:1000-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.4-0.5 mg/mL BSA, 0.02% sodium azide).
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of (1) A431 cell lysate, (2) A431 lysate cell treated with EGF using STAT1 (phospho Y701) monoclonal antibody, clone HHB-19.

Gene Info — STAT1

Entrez GeneID [6772](#)

Protein Accession# [P42224](#)

Gene Name STAT1

Gene Alias DKFZp686B04100, ISGF-3, STAT91

Gene Description signal transducer and activator of transcription 1, 91kDa

Omim ID [209950 600555](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]

Other Designations OTTHUMP00000165047|signal transducer and activator of transcription 1|signal transducer and activator of transcription-1|transcription factor ISGF-3 components p91/p84

Pathway

- [Chemokine signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)

- [Pathways in cancer](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Campylobacter Infections](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Glomerulonephritis](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)
- [Infant](#)

- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neutropenia](#)
- [Osteoporosis](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Proteinuria](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Salmonella Infections](#)
- [Thrombocytopenia](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Werner syndrome](#)