


PIAS1 monoclonal antibody, clone ABFA-16

Catalog # MAB20594

Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of HepG2 cell lysate using PIAS1 monoclonal antibody, clone ABFA-16.

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human PIAS1.

Immunogen A synthetic peptide corresponding to human PIAS1.

Host Rabbit

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage

- Flow Cytometry (1:50)
- Immunocytochemistry (1:50-1:200)
- Immunofluorescence (1:50-1:200)
- Immunohistochemistry (1:50-1:200)
- Immunoprecipitation (1:50)
- Western Blot (1:500-1:2000)
- The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).

Storage Instruction

Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of HepG2 cell lysate using PIAS1 monoclonal antibody, clone ABFA-16.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

- Immunocytochemistry

- Immunofluorescence

- Immunoprecipitation

- Flow Cytometry

Gene Info — PIAS1

Entrez GeneID[8554](#)**Protein Accession#**[O75925](#)**Gene Name**

PIAS1

Gene Alias

DDXBP1, GBP, GU/RH-II, MGC141878, MGC141879, ZMIZ3

Gene Description

protein inhibitor of activated STAT, 1

Omim ID[603566](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a member of the mammalian PIAS [protein inhibitor of activated STAT-1 (signal transducer and activator of transcription-1)] family. This member contains a putative zinc-binding motif and a highly acidic region. It inhibits STAT1-mediated gene activation and the DNA binding activity, binds to Gu protein/RNA helicase II/DEAD box polypeptide 21, and interacts with androgen receptor (AR). It functions in testis as a nuclear receptor transcriptional coregulator and may have a role in AR initiation and maintenance of spermatogenesis. [provided by RefSeq]

Other Designations

AR interacting protein|DEAD/H (Asp-Glu-Ala-Asp/His) box binding protein 1|protein inhibitor of activated STAT-1|zinc finger, MIZ-type containing 3

Pathway

- [Jak-STAT signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [Ubiquitin mediated proteolysis](#)

Disease

- [Multiple Sclerosis](#)