


TRAF2 monoclonal antibody, clone HOG-20

Catalog # MAB20801 Size 100 uL

Applications



Western Blot (Cell lysate)

Western Blot analysis of HeLa cell lysate with TRAF2 monoclonal antibody, clone HOG-20 (Cat # MAB20801).

Specification

Product Description Rabbit monoclonal antibody raised against synthetic peptide of human TRAF2.

Immunogen A synthetic peptide corresponding to human TRAF2.

Host Rabbit

Theoretical MW (kDa) 55.859

Reactivity Human

Form Liquid

Purification Affinity purification

Isotype IgG

Recommend Usage

- Flow Cytometry (1:50)
- Immunocytochemistry (1:50-1:200)
- Immunofluorescence (1:50-1:200)
- Immunohistochemistry
- Immunoprecipitation (1:40)
- Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide).
Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western Blot analysis of HeLa cell lysate with TRAF2 monoclonal antibody, clone HOG-20 (Cat # MAB20801).

- Immunohistochemistry

- Immunocytochemistry

- Immunofluorescence

- Immunoprecipitation

- Flow Cytometry

Gene Info — TRAF2

Entrez GeneID	7186
Protein Accession#	Q12933
Gene Name	TRAF2
Gene Alias	MGC:45012, TRAP, TRAP3
Gene Description	TNF receptor-associated factor 2
Omim ID	601895
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can ubiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq]

Other Designations

OTTHUMP00000022625|OTTHUMP00000064745|tumor necrosis factor type 2 receptor associated protein 3

Pathway

- [Adipocytokine signaling pathway](#)
- [Apoptosis](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)

Disease

- [Alzheimer disease](#)
- [Cardiovascular Diseases](#)
- [Connective Tissue Diseases](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Ductus Arteriosus](#)
- [Edema](#)
- [Fetal Diseases](#)
- [Genetic Predisposition to Disease](#)

- [Hematologic Diseases](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Retinopathy of Prematurity](#)
- [Skin Diseases](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)